



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Mathematics and Computers in Simulation xxx (2006) xxx–xxx

www.elsevier.com/locate/matcom

Computing the principal eigenvalue of the Laplace operator by a stochastic method

Antoine Lejay^{a,*}, Sylvain Maire^b

^a *Projet OMEGA (INRIA/IECN), IECN, Campus Scientifique, BP 239, 54506 Vandoeuvre-lès-Nancy Cedex, France*

^b *ISITV, Université de Toulon et du Var, Avenue G. Pompidou, BP 56, 83262 La Valette du Var Cedex, France*

Received 6 December 2005; accepted 8 June 2006

Abstract

We describe a Monte Carlo method for the numerical computation of the principal eigenvalue of the Laplace operator in a bounded domain with Dirichlet conditions. It is based on the estimation of the speed of absorption of the Brownian motion by the boundary of the domain. Various tools of statistical estimation and different simulation schemes are developed to optimize the method. Numerical examples are studied to check the accuracy and the robustness of our approach.

© 2006 IMACS. Published by Elsevier B.V. All rights reserved.

MSC: 65C05; 60F15

Keywords: First eigenvalue of the Dirichlet problem; Euler scheme for Brownian motion; Random walk on spheres; Random walk on rectangles

1. Introduction

The aim of this paper is to introduce and study a Monte Carlo method for the numerical computation of the principal eigenvalue of the Laplace operator $\frac{1}{2}\Delta$ in a bounded domain $D \subset \mathbb{R}^d$ with a sufficiently piecewise smooth boundary ∂D and with Dirichlet boundary conditions. This leading eigenvalue determines the speed of convergence to the steady state for the solution of the heat equation. To compute it by a deterministic method, one has to discretize the Laplace operator using for example finite differences or finite elements and then evaluate the largest eigenvalue of the discretization matrix using for example the inverse power method. This computation becomes really expensive when the spatial dimension d increases. Moreover the discretization should be refined enough so that the principal eigenvalue of the discretization matrix is close enough to the principal eigenvalue of the Laplace operator. Hence it is worth considering Monte Carlo methods [15] because they are usually efficient for this kind of difficult problems since they do not necessarily require to discretize the domain D and they depend only linearly on the spatial dimension.

We have introduced in [17] a stochastic method to compute the principal eigenvalue of neutron transport operators based on the numerical computation of the type of the neutron transport operator. The idea was to combine the formal eigenfunction expansion of the solution of the relative Cauchy problem and its Monte Carlo evaluation via the

* Corresponding author. Tel.: +33 3 83 68 45 51; fax: + 33 3 83 68 45 34.

E-mail addresses: Antoine.Lejay@iecn.u-nancy.fr (A. Lejay), maire@univ-tln.fr (S. Maire).

Feynman–Kac formula. We intend to use the same methodology here in the case of the Laplace operator. The stochastic representation of the principal eigenvalue of the operator $\frac{1}{2}\Delta$ is usually achieved by combining the solution $u(t, x)$ of the Cauchy problem

$$\frac{\partial u}{\partial t} = \frac{1}{2}\Delta u, \quad u(0, x) \equiv 1, \quad x \in D \subset \mathbb{R}^d$$

obtained by the Feynman–Kac formula and the formal eigenfunction expansion

$$u(t, x) = \sum_{j=1}^{\infty} c_j \exp(\lambda_j t) \Psi_j(x)$$

of this solution in $L^2(D)$ where λ_j are the eigenvalues of $\frac{1}{2}\Delta$ arranged in decreasing order and Ψ_j their relative eigenfunctions. Indeed as the solution of this equation is given by

$$u(t, x) = \mathbb{P}(\tau_D^x > t),$$

where τ_D^x is the exit time from D of the Brownian motion starting at x , the principal eigenvalue λ_1 is directly linked to the speed of absorption of the Brownian motion by the boundary ∂D and we have for all $x \in D$

$$\lambda_1 = \lim_{t \rightarrow +\infty} \frac{1}{t} \log \mathbb{P}(\tau_D^x > t).$$

This result is also true for a general elliptic operator A with Dirichlet boundary conditions in a bounded domain D , where τ_D^x is the exit time from D of the stochastic process X_t^x generated by A [7,8,13,19]. We had to compute numerically the same quantity in [17] when studying homogeneous neutron transport operator. In this case, an exact simulation of transport processes involved in Feynman–Kac representations was possible. In the case of the Brownian motion, one has to use approximations based on various discretization schemes.

In Section 2, we describe quickly grid-free schemes, some of them common and some of them new, that can be used here. Then, we present in Section 3 different estimators for λ_1 based on an accurate study of the eigenfunctions expansion of the solution $u(t, x)$. Some of these estimators were developed in [17] but we also introduced new ones based on correlation coefficients. In order to compare the different simulations schemes and the different estimators, we study in detail in Section 4 a bidimensional problem. We finally study in Section 5 more difficult problems in dimensions 3 and 5 combining all the tools developed previously in order to show that our method is also relevant in these cases.

2. Exit time simulation procedures

The aim of this section is to describe and compare some simulation schemes for the exit time of the Brownian motion in a bounded domain D with Dirichlet boundary conditions.

2.1. Euler schemes

The Euler scheme (see for example [14]) with discretization parameter Δt writes

$$X_0 = x, \quad X_{n+1} = X_n + \sqrt{\Delta t} Y_n$$

where the Y_n are independent standard Gaussian random variables. To compute $\mathbb{P}(\tau_D^x > t)$ one needs to compute simulations of τ_D^x . With the crude version, the simulation stops once $X_{n+1} \in D^c$ and τ_D^x is approximated either by $n\Delta t$, $(n + \frac{1}{2})\Delta t$ or a slightly refined approximation based on the distances $d_n = d(X_n, \partial D)$ and $d_{n+1} = d(X_{n+1}, \partial D)$. In any case, these approximations are of weak order $\sqrt{\Delta t}$ and they overestimate this exit time. Indeed, the main simulation error does not come from the error at the last step but from the possibility for the Brownian motion to leave the domain between step n and $n + 1$ and be back into it at time $(n + 1)\Delta t$. It is possible to take into account this possibility to obtain a scheme of weak order Δt using the half-space approximation [9,10]. An additional random test based on d_n

and d_{n+1} is required. Taking a uniform random variable U_n , the motion stops if

$$\exp\left(-\frac{2d_n d_{n+1}}{\Delta t}\right) > U_n.$$

It is still possible to introduce another refinement at the last step [3] which improves the accuracy but does not change the order of the scheme. In [11], another method is proposed that leads to the same order of convergence as in the half-space method. We only consider in the examples the naive version and the one using the half-space approximation, which we denote from now by the Improved Euler scheme.

2.2. Walk on spheres (WOS)

The simulation method based on the Euler schemes covers a wide range of elliptic and parabolic partial differential equations and can take into account the dependence of the drift and of the diffusion coefficients on the spatial variables. One only has to simulate a stochastic differential equation by means of this scheme instead of the Brownian motion. However, in the case of the heat equation some faster schemes are available. The walk on sphere schemes (WOS) [20] relies on the isotropy of the Brownian motion. This walk goes from x to the boundary ∂D from a sphere to another until the motion reaches an ε -absorption layer. The spheres are built so that the jumps are as large as possible. The radius of the next sphere from a starting point x_n is $d(x_n, \partial D)$ and the next point is chosen uniformly on this sphere. The average number of steps to exit from the domain is proportional to $|\log(\varepsilon)|$ [20]. As we want here to simulate τ_D^x , we need in addition to simulate the law $Y(r)$ of the exit time from a sphere of radius r starting at its center. Some scaling arguments on the partial differential equation or on the Brownian motion show that $r^2 Y(1)$ and $Y(r)$ have the same law. Some analytical expressions for the distribution function $H(t) = \mathbb{P}(Y(1) < t)$ are available. In dimension 2, we have for example [1]

$$H(t) = \sum_{k=1}^{\infty} \frac{2J_0(0)}{j_k J_1(j_k)} \exp\left(\frac{-j_k^2 t}{2}\right)$$

where J_0 and J_1 are Bessel functions and the j_k are the positive zeros of J_0 . The simulation of $Y(1)$ by the standard method requires the inversion of a series which can be difficult and costly. As we only need here to simulate $Y(1)$ during all the walk, we tabulate the distribution function once and for all and use it for simulations. Note that we can either use its analytical expression to do it or Monte Carlo simulations based on the corrected Euler scheme with a small parameter and a huge number of simulations. The second method is especially efficient in higher dimensions.

Some refined versions (for example, the walk on spheres with shifted centers [12]) allow faster absorption by the boundary, but also leads to simulate complex random variables.

2.3. Walk on rectangles (WOR)

In many practical situations (temperature evolution in a room, . . .), the boundary is or can be approximated by a polygon. In this case, exact and fast simulations are possible.

The idea of the random walk on rectangles (WOR) is a generalization of the random walk on squares [4], that can be deduce from the algorithm given in [18] to simulate stochastic differential equations (SDE). The idea is to simulate the exit time and position from a rectangle (or a parallelepiped in dimension greater than 2) by a Brownian motion. Unlike with a sphere, the random variables giving the exit time and position from a rectangle are not independent. Yet, as shown in [5], this could be achieved rather efficiently by proper conditioning and reducing the problem to simulate random variables related to the 1D Brownian motion. The advantage of this method is that, unlike with the random walks of spheres and squares, the rectangles may be chosen prior to any simulation, since the Brownian motion can start from any point in it. In addition, it can be used even if a constant drift term is present, or to deal with Neumann boundary condition. Although it takes more time to simulate the exit time and position from a rectangle, the number of simulations is reduced. Of course, one has to use, when possible, rectangles for which at least one side corresponds to a boundary of D .

Let (B^1, \dots, B^d) be a d -dimensional Brownian motion with the starting point $(B_0^1, \dots, B_0^d) = (x^1, \dots, x^d)$. We denote by \mathbb{P}_x the distribution of the 1D Brownian motion starting at x . We are interested in simulating its first exit time

and position from the parallelepiped $[-L_1, L_1] \times \cdots \times [-L_d, L_d]$. For that, we set $\tau^i = \inf\{t > 0; B_t^i \notin [-L_i, L_i]\}$ for $i = 1, \dots, d$ and we perform roughly the following operations:

- (1) We draw a realization (θ^1, y^1) for $(\tau^1, B_{\tau^1}^1)$ under \mathbb{P}_{x^1} . We set $\bar{\tau} = \theta_1, S_1 = 1$ and $J = 1$.
- (2) For i from 2 to d , do
 - (2.1) We use a Bernoulli random variable to decide whether $\{\tau^i < \bar{\tau}\}$ or $\{\tau^i \geq \bar{\tau}\}$ under \mathbb{P}_{x^i} .
 - (2.2) If we have decided that $\{\tau^i < \bar{\tau}\}$, then we draw a realization (θ^i, y^i) of $(\tau^i, B_{\tau^i}^i)$ under \mathbb{P}_{x^i} given $\{\tau^i < \bar{\tau}\}$. We set $J = i, S_i = 1$ and $\bar{\tau} = \tau^i$.
 - (2.3) Otherwise, we set $S_i = 0, \varphi_i = \bar{\tau}$.
- (3) We set $z^J = y^J$.
- (4) For i from 1 to d , do
 - (4.1) If $S_i = 1$ and $i \neq J$ then we simulate a realization z^i of $B_{\bar{\tau}}^i$ under \mathbb{P}_{x^i} given $\{\tau^i = \theta^i, B_{\tau^i}^i = y^i\}$.
 - (4.2) Otherwise, if $S_i = 0$, then we simulate a realization z^i of $B_{\varphi_i}^i$ under \mathbb{P}_{x^i} given $\{\tau^i \geq \varphi_i\}$.
- (5) The algorithm stops and returns the time $\bar{\tau}$ and the position (z^1, \dots, z^d) on the boundary of the parallelepiped.

The involved distributions and the exact algorithm are detailed in [5].

2.4. Walk on rectangles with importance sampling (WOR-IS)

This method, developed in [6], is a variation on the previous method. Instead of simulating the exact couple exit time τ and exit position B_τ of a rectangle R for a Brownian motion B , we draw this exit time and position (θ, Z) using an arbitrary distribution (of course, chosen to be simple), and we compute a weight w such that $\mathbb{E}(f(\tau, B_\tau)) = \mathbb{E}(wf(\theta, Z))$ for any bounded, measurable function f defined on $\mathbb{R}_+ \times \partial R$. Of course, for any function g defined on $\mathbb{R}_+ \times \partial D$, the functional $\mathbb{E}(g(\tau_D^x, B_{\tau_D^x}))$ is then evaluated by $\mathbb{E}(w_1 \dots w_{n^*} g(\theta_{n^*}, Z_{n^*}))$, where the (θ_i, Z_i) 's are the successive exit times and positions for a sequence of rectangles, the w_i 's are their associated weights, and n^* is the first integer for which Z_{n^*} belongs to ∂D , or is close enough to ∂D .

The advantage of this method over the WOR/WOS is that we are free to choose the distribution of (θ, Z) , and then we can easily “constraint” the diffusion to go in some direction or condition it not to reach a part of the boundary of the domain. Thus, this method can be used to simulate Brownian motion or SDE in domains with complex geometries and to reduce the variance of the estimators that are computed. Moreover, this method is much more faster than the WOR if we choose for (θ, Z) random variables that are easy to simulate. In addition, the weights are rather easily computed since they rely on the density of the first exit time and the position of a killed Brownian motion in the 1D case.

However, this method suffers from a severe drawback in our case. The empirical distribution function $F_N(t)$ of $F(t) = \mathbb{P}(\tau_D^x < t)$ is constructed as follows. If $(\theta^{(i)}, w^{(i)})_{i=1, \dots, N}$ are the simulated exit time from D with their associated weights $w^{(i)}$, then

$$F_N(t) = \frac{1}{N} \sum_{i=1}^N w^{(i)} \chi_{[0, t)}(\theta^{(i)}).$$

If $\hat{F}_N(t)$ is the empirical distribution function constructed with N independent random variables with the same law as τ_D^x , then $\text{Var}(\hat{F}_N(t)) = N^{-1} F(t)(1 - F(t))$. In our case,

$$\text{Var}(F_N(t)) = \frac{1}{N} \text{Var}(w^{(1)} \chi_{[0, t)}(\theta^{(1)})).$$

The problem is then to find a “strategy” for which $\text{Var}(F_N(t))$ is smaller than $\text{Var}(\hat{F}_N(t))$, at least when t is large enough. Unfortunately, we have not been able to find a good way to do so. For the sake of simplicity, we have adopted the following strategy. For each rectangle, each of its side is reached with a probability 1/4, the exit position is drawn

using a uniform random variable on the side and the exit time is simulated using an exponential random variable of parameter 0.35 times the square of the length of the side that is not reached.

3. Estimators of the principal eigenvalue

3.1. Direct approximation

Using the formal eigenfunction expansion of $u(x, t)$, we can write that almost everywhere

$$\frac{1}{t} \log(\mathbb{P}(\tau_D^x > t)) = \frac{1}{t} \log(C_1 e^{\lambda_1 t} + C_2 e^{\lambda_2 t} + \dots + C_k e^{\lambda_k t} + o(e^{\lambda_k t})),$$

where $C_1 = c_1 \Psi_1(x)$. The Krein–Rutman theorem [19] ensures that λ_1 is simple and $C_1 > 0$. Then

$$\frac{1}{t} \log \mathbb{P}(\tau_D^x > t) \simeq \lambda_1 + \frac{\log(C_1)}{t} + \frac{C_2}{C_1} \frac{e^{(\lambda_2 - \lambda_1)t}}{t}$$

keeping only the two dominant terms. The previous approximation shows that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(\tau_D^x > t) = \lambda_1$$

almost everywhere and it has been proved in [13] that it holds everywhere. Hence the Monte Carlo computation of $t^{-1} \log \mathbb{P}(\tau_D^x > t)$ for large times offers a first possibility to give a numerical approximation of λ_1 . Moreover we can choose the starting point wherever we want. The most natural choice is to take this starting point in the center of the domain in order to make the trajectories last as long as possible. Even if we do so, we will see in some numerical examples that this direct method gives in fact a poor approximation because the variance increases quickly with time. Indeed this involves the computation of the probability of rare events which become rarer and rarer as time increases. We now give some better estimators to overcome this difficulty.

3.2. Interpolation method

The previous expansion is constituted of a main term $\lambda_1 + t^{-1} \log(C_1)$ and of a term $\frac{C_2}{C_1} \frac{e^{(\lambda_2 - \lambda_1)t}}{t}$ which decays at an exponential rate the more quickly the first two eigenvalues are distant from each other. The difference $\lambda_2 - \lambda_1$ is intrinsic to the domain. Nevertheless the choice of a starting point near the center of the domain makes the ratio C_2/C_1 smaller. This can be seen for instance on the eigenfunctions expansion on square domains. Moreover, it is well known that the errors due to the discretization scheme are smaller for points away from the boundary. If we assume that t is large enough so that this term is negligible with respect to the others, we can compute easily λ_1 using the Monte Carlo approximations \bar{p}_1 and \bar{p}_2 of, respectively, $p_1 = \mathbb{P}(\tau_D^{x,S} > t_1)$ and $p_2 = \mathbb{P}(\tau_D^{x,S} > t_2)$ with a discretization scheme S . We obtain the estimators

$$\lambda^{\text{In}}(t_1, t_2) = \frac{\log(\bar{p}_2) - \log(\bar{p}_1)}{t_2 - t_1}$$

of λ_1 which are also used in neutron transport criticality computations [2]. Note that \bar{p}_1 and \bar{p}_2 are obviously correlated and may also be biased estimators of $\mathbb{P}(\tau_D^x > t_1)$ and $\mathbb{P}(\tau_D^x > t_2)$ because of the discretization errors. Letting $s_1 = \sqrt{\bar{p}_1(1 - \bar{p}_1)}$, we obtain the 95% confidence interval

$$\bar{p}_1 - 1.96 \frac{s_1}{\sqrt{N}} \leq p_1 \leq \bar{p}_1 + 1.96 \frac{s_1}{\sqrt{N}}$$

that is assuming that $\frac{s_1}{\bar{p}_1 \sqrt{N}}$ is small enough,

$$\log(\bar{p}_1) - 1.96 \frac{s_1}{\bar{p}_1 \sqrt{N}} \leq \log(p_1) \leq \log(\bar{p}_1) + 1.96 \frac{s_1}{\bar{p}_1 \sqrt{N}}.$$

We finally have the 90% confidence interval for $\lambda^{\text{In}}(t_1, t_2)$ writing

$$\frac{\log\left(\frac{\bar{p}_2}{\bar{p}_1}\right)}{t_2 - t_1} - \frac{1.96N^{-1/2}}{t_2 - t_1} \left(\frac{s_1}{\bar{p}_1} + \frac{s_2}{\bar{p}_2}\right) \leq \lambda^{\text{In}}(t_1, t_2) \leq \frac{\log\left(\frac{\bar{p}_2}{\bar{p}_1}\right)}{t_2 - t_1} + \frac{1.96N^{-1/2}}{t_2 - t_1} \left(\frac{s_1}{\bar{p}_1} + \frac{s_2}{\bar{p}_2}\right). \quad (1)$$

3.3. Least squares approximations

In order to make a global use of the information given by the computation of the solution at the different times, we can give a least squares approximation of λ_1 and $\log(K_0)$ by fitting these parameters to the approximation model still assuming that the term $\frac{C_2}{C_1} \frac{e^{(\lambda_2 - \lambda_1)t}}{t}$ is negligible. We denote by $F_N(t)$ be the empirical distribution of τ_D^x , which is known at the set of times t_i . We have to find λ_1 and $\beta = \log(C_1)$ minimizing

$$\sum_{i=p}^q \left(\lambda_1 + \frac{\beta}{t_i} - \frac{1}{t_i} \log(1 - F_N(t_i)) \right)^2,$$

which is a linear least squares problem with respect to the parameters λ_1 and β . This method was tested successfully in [16,17] but we adopt a slightly different approach which appears to be more accurate and rigorous.

We now let $F(t) = \mathbb{P}(\tau_D^x < t)$ be the distribution function of the exit time for a fixed point x . Instead of computing λ_1 as the intersection value in the linear regression, we compute it as the slope of the regression line, since

$$\log(1 - F(t)) = \lambda_1 t + \log(c_1 \Psi_1(x)) + \varepsilon(t)$$

with

$$\varepsilon = \log(1 + R(t, x)/c_1 \Psi_1(x)) \quad \text{and} \quad R(t, x) = \sum_{k \geq 2} e^{-(\lambda_k - \lambda_1)t} \Psi_k(x) c_k.$$

Of course, in addition to ε , a second error comes from the replacement of $F(t)$ by an empirical density $F_N(t)$, where N is the number of particles in the simulation. Thus, we are indeed using a linear regression for the equation

$$\log(1 - F_N(t)) = -\lambda_1 t + \log(\alpha_1 \Psi_1(x)) + \varepsilon(t) + \eta_N(t)$$

with

$$\eta_N(t) = \log\left(1 - \frac{F(t) - F_N(t)}{1 - F(t)}\right) \approx \frac{F_N(t) - F(t)}{1 - F(t)}.$$

The speed of convergence of $F_N(t)$ to $F(t)$ given by the Kolmogorov–Smirnov test [21] is of order $1/\sqrt{N}$. We will see that our estimate of λ_1 is more likely to be limited by η_N , and thus by the number N of samples than by ε .

We can be even more precise. The empirical distribution function $F_N(t)$ converges uniformly to $F(t)$. Moreover, $\sqrt{N}(F_N(t) - F(t))$ converges in distribution to a Gaussian process $(x_t)_{t \geq 0}$ with $\text{Var}(x_t) = F(t)(1 - F(t))$ and $\text{Cov}(x_t, x_s) = F(s)(1 - F(s))$ when $t \geq s$. This implies that

$$\text{Var}(\eta_N(t)) \approx \frac{1}{N} \frac{F(t)}{1 - F(t)},$$

which was already used in the interpolation method. Thus, a good estimate of λ_1 relies on using a window in which both $\varepsilon(t)$ (which is unknown) and $\text{Var}(\eta_N(t))$ are small enough.

Remark 1. We could also use weighted least squares methods with weights given by $(1 - F_N(t))/F_N(t)$ in order to compensate the variance of the $\eta_N(t)$. In practice, we have noticed no real improvement.

4. A 2D test case

4.1. Description

We deal with the 2D domain seen in Fig. 1




Fig. 1. The domain D for the 2D test case.

First, we estimate the first eigenvalue λ_{2D} of $\frac{1}{2}\Delta$ using the `pdetool` package from MATLAB. In Table 1, we give the estimates of λ_{2D} as a function of the number of times we refine the mesh. Thus, it seems that the real value of λ_{2D} is close to -0.740 . The order of the second eigenvalue is -1.75 .

4.2. The empirical distribution function of the first exit time by Monte Carlo methods

We have considered the simulation of the function $F(t)$ using the different schemes described in Section 2 with 1,000,000 particles, excepted for the WOR-IS where 100,000,000 particles were used (see Section 2.4 for an explanation). For the Euler scheme, we use a time step Δt of 10^{-3} . For the walk on spheres, the absorption boundary layer is $\varepsilon = 10^{-3}$. For the WOR and WOR-IS, we use only the two rectangles $[0, 4] \times [0, 3]$ and $[1, 3] \times [0, 4]$ (the lower left corner is at $(0,0)$). The computation times are presented in Section 4.5.

In Fig. 2, we have drawn $G_N(t) = t^{-1} \log(1 - F_N(t))$ where the starting point is $(2, 2)$. All the schemes give a function G_N decreasing to a value which is around -0.7 at $t = 10$, except for the naive Euler scheme for which it is around -0.67 . For the WOR-IS, we have tested different strategies to force the process to stay longer times in the domain, but none were very successful. In addition, we see that the curves are perturbed by numerical artifacts.

Yet for large times, that is $t > 7.0$, the behavior becomes quite erratic, since it corresponds to the simulation of rare events. We cannot expect to obtain an accurate estimate of λ_{2D} this way.

4.3. Estimation by the interpolation method

We then use the interpolation method of Section 3.2, with $t_2 - t_1 = 4$ and we show in Fig. 3 the values of $\lambda_1^S(t_1, t_2)$ for $t_1 \in [2, 5]$.

We obtain more accurate estimators than with the previous method. We still observe a deterioration when t_1 is too large for all the schemes. However, the naive Euler scheme gives an overestimated value of λ_{2D} , which can be easily explained by its nature. We no longer consider this scheme in the following.

If we look for the value of $\lambda^{\text{In}}(t_1, t_1 + 4)$ with t_1 in some particular intervals, it e.g. $t_1 \in [2, 3]$, we can see that all the estimators are very close to -0.74 and this for all the remaining schemes. We now turn to more quantitative results.

Table 1
Estimate by MATLAB/`pdetool` in function of the refinement of the mesh

Number of nodes	$-\lambda_{2D}$
180	0.7578
670	0.74532
2590	0.74415
10200	0.74021
40400	0.73975
160940	0.73958




Fig. 2. $t^{-1} \log(1 - F_N(t))$ for the different schemes.

4.4. Optimization of the estimators

To estimate λ_{2D} , we are free to choose the range of time t for $F(t)$. The idea is then to consider an estimator $\lambda_{2D}^{(i)}$ of λ_{2D} using the data on an interval $[a_i, b_i]$ for a large choice of a_i and b_i . We perform a statistical analysis of the set of the $\lambda_{2D}^{(i)}$ for which a quality criterion is satisfied.

We first use the confidence interval techniques of Section 3.2 after having remarked in Fig. 3 that all the estimators $\lambda^{\text{In}}(t_1, t_1 + 4)$ are contained in a narrow band for $t_1 \in [2, 5]$. We now compute all the possible estimators $\lambda^{\text{In}}(t_1, t_2)$ and their relative confidence intervals, where t_1, t_2 are chosen on a grid $\eta\mathbb{Z}$ with $\eta = 1/10$, $t_1 \geq 2$, $t_2 \leq 8$ and $t_2 - t_1 \geq 2$. We only keep the fraction of these estimators for which the length of the confidence interval is small. Indeed, we have to find a balance in the length $\ell_{90\%}(t_1, t_2)$ of the confidence interval at 90% given by (1) between the term




Fig. 3. The estimator $\lambda^{\text{In}}(t_1, t_1 + 4)$ for the different schemes.




Fig. 4. $\lambda^{\ln}(t_1, t_2)$ in function of $\ell_{90\%}(t_1, t_2)$ for the improved Euler scheme.

$1/(t_2 - t_1)$ which shall be small and the term $s_1/\bar{p}_1 + s_2/\bar{p}_2$ that increases when t_1 or t_2 increases. We plot $\lambda^{\ln}(t_1, t_2)$ against $\ell_{90\%}(t_1, t_2)$ and we remark that the quality of the approximation decreases when $\ell_{90\%}(t_1, t_2)$ increases (Fig. 4).

We also note that the smallest values of $\ell_{90\%}(t_1, t_2)$ are obtained when t_1 is close to 2 and t_2 close to 5.

We now present our results in Table 2 using classical statistical estimators for the $\lambda^{\ln}(t_1, t_2)$'s satisfying our criterion.

If we look at the median or the mean of the estimators $\lambda^{\ln}(t_1, t_2)$, we can reach an accuracy of at least 3×10^{-3} . All the schemes give an analogous accuracy.

We now turn to the least squares method, by replacing our criterion on the length of the confidence interval by a criterion based on the correlation coefficient R . This coefficient measures the validity of the linear approximation. We

Table 2

Study of the estimators $\lambda^{\ln}(t_1, t_2)$ by keeping the 10% having the smallest value of $\ell_{90\%}(t_1, t_2)$

	Method			
	Improved Euler	WOS	WOR	WOS-IS
Minimum	-0.7423	-0.7385	-0.7398	-0.7537
1st Qu.	-0.7403	-0.7378	-0.7376	-0.7456
Median	-0.7392	-0.7373	-0.7373	-0.7393
Mean	-0.7391	-0.7372	-0.7374	-0.7404
3rd Qu.	-0.7377	-0.7368	-0.7369	-0.7364
Maximum	-0.7365	-0.7351	-0.7361	-0.7268

Table 3

Study of the estimators $\lambda^{\text{LS}}(t_1, t_2)$ in function of R^2 for the improved Euler scheme

	Method		
	Improved Euler	WOS	WOR
Minimum	-0.7470	-0.7415	-0.7443
1st Qu.	-0.7444	-0.7375	-0.7411
Median	-0.7413	-0.7368	-0.7396
Mean	-0.7423	-0.7367	-0.7398
3rd Qu.	-0.7401	-0.7360	-0.7380
Maximum	-0.7395	-0.7321	-0.7370




Fig. 5. $\lambda^{\text{LS}}(t_1, t_2)$ in function of R^2 for the improved Euler scheme.

denote by $\lambda^{\text{LS}}(t_1, t_2)$ the estimator of $\lambda^{\text{LS}}(t_1, t_2)$ giving the slope of the linear regression of $G_N(t)$ with $t \in [t_1, t_2]$. We select the estimators $\lambda^{\text{LS}}(t_1, t_2)$ for which R^2 is greater than 0.99997. We also plot $\lambda^{\text{LS}}(t_1, t_2)$ against the coefficient R^2 . We now present our results on $\lambda^{\text{LS}}(t_1, t_2)$ in Table 3 as we did previously for $\lambda^{\text{In}}(t_1, t_2)$ (Fig. 5).

Once again, we obtained a good accuracy on the first eigenvalue λ_{2D} (at least 3×10^{-3}). However, for the WOR-IS, this criterion is less robust and we have obtained an accuracy 3×10^{-2} and thus we do not include the results in Table 3.

4.5. Computation times

Although the schemes give close estimated values of λ_{2D} , their relative computation times differ largely and may be function of the choice of some parameters, as seen in Table 4. The simulations were done on a bi-processors DEC 700 MHz.

We compare first the average number of steps of each of the methods. The WOR and WOR-IS take around two steps to reach the boundary. The WOS takes around ten times more steps using a boundary layer $\varepsilon = 10^{-4}$. This number of steps increases slowly like a $O(|\log(\varepsilon)|)$ as ε goes to zero. The Euler scheme with $\Delta t = 10^{-2}$ takes 10 times more steps than the WOS. This number of steps increases linearly with Δt (if Δt is divided by 10 the number of steps is ten times greater). If we now look at the CPU times, the WOR-IS is really the fastest method as its average number of steps is 2 and that it requires the simulation of fairly simple random variables. Unfortunately, this method increases somehow the variance of the simulations. A lot more samples are required to make it as accurate as the other methods. The simulation times of the WOR and of the Euler scheme are comparable. The first one requires only 1.5 steps but the simulation of complex random variables. The second one requires many steps but the simulation of very simple

Table 4
Computations times with $N = 1,000,000$ for all the methods

Scheme	Parameter name	Parameter	Time (s)	Average number of steps
Euler	Time step, Δt	10^{-2}	210	200
		10^{-3}	1400	2000
WOS	Boundary layer, ε	10^{-3}	80	13
		10^{-4}	85	16
WOR	–	–	700	1.5
WOR-IS	–	–	16	2.1

Gaussian random variables. A good arrangement between these methods appears to be the WOS. The average number of steps is small and the random variables to simulate are not too complex.

4.6. Preliminary conclusions

Our approach is efficient on this test case. However, among the schemes we have tested, the naive Euler scheme shall be discarded. Our statistical study gives an approximation of λ_{2D} with an accuracy of order 3×10^{-3} , which corresponds to what can be expected from the replacement of $F(t)$ by an empirical distribution function, due to the Kolmogorov–Smirnov theorem.

In practice, we recommend the two following methods:

- We look graphically at either $\lambda^{\text{In}}(t_1, t_1 + \tau)$ for a given τ and t_1 in the range of the obtained values and we then determine a range of times for which the oscillations around a constant value are small enough. We may use either the least squares method or the interpolation method.
- We use the least squares methods to estimate $\lambda^{\text{LS}}(t_1, t_2)$ for a large number of values of $t_1 < t_2$ with $t_2 - t_1 \geq \tau$ for an arbitrary τ , and keep the values of $\lambda^{\text{LS}}(t_1, t_2)$ for which the coefficient R^2 is large enough.

5. More complex problems

5.1. A 3D problem

We now consider the extension of our 2D problem where we add a third dimension $z \in [0, 4]$. We could treat as well more complex three dimensional domains but we use this problem as its main eigenvalue can be simply computed. By a separation of variables argument, we have $\lambda_{2D} - \frac{\pi^2}{32} \simeq -1.048$.

In Table 5, we use confidence intervals and least squares approximations for intervals $[t_1, t_2]$ contained in $[2, 8]$ with $t_2 - t_1 \geq 2$ and a step of 0.1 for t_1 and t_2 .

We focus on this example on the WOR and WOS methods. For the WOS, we need to simulate the uniform law on the unit sphere. This density writes

$$f(\theta, \varphi) = \frac{1}{2} \sin(\theta) \mathbf{1}_{[0, \pi)}(\theta) \frac{1}{2\pi} \mathbf{1}_{[0, 2\pi)}(\varphi).$$

The three Cartesian coordinates are

$$x = \sin(\theta) \sin(\varphi), \quad y = \sin(\theta) \cos(\varphi), \quad z = \cos(\theta)$$

Table 5

Study of the estimators $\lambda^{\text{In}}(t_1, t_2)$ for $\ell_{90\%}(t_1, t_2) \leq \ell_{\max}$ and $\lambda^{\text{LS}}(t_1, t_2)$ for $R^2 \geq R_{\min}^2$ with $N = 1,000,000$ particles

	Method			
	WOR		WOS	
Parameter, ε	–		10^{-4}	
Time (s)	1200		110	
# steps	1.34		30	
	WOR		WOS	
	λ^{LS}	λ^{In}	λ^{LS}	λ^{In}
R_{\min}^2 or ℓ_{\max}	0.99995	0.02	0.99995	0.02
Number of samples	141	125	695	135
Minimum	–1.056	–1.048	–1.049	–1.049
1st Qu.	–1.049	–1.045	–1.047	–1.048
Median	–1.046	–1.044	–1.047	–1.047
Mean	–1.047	–1.044	–1.046	–1.047
3rd Qu.	–1.045	–1.043	–1.046	–1.046
Maximum	–1.041	–1.042	–1.041	–1.045

Table 6

Study of the estimators $\lambda^{\text{In}}(t_1, t_2)$ for $\ell_{90\%}(t_1, t_2) \leq \ell_{\max}$ and $\lambda^{\text{LS}}(t_1, t_2)$ for $R^2 \geq R_{\min}^2$ with $N = 1,000,000$ particles

Parameter, Δt	Method					
	WOR		Improved Euler		Improved Euler	
	2000		300		440	
Time (s)	1.23		123		200	
Number of steps	WOR		Improved Euler		Improved Euler	
	λ^{LS}	λ^{In}	λ^{LS}	λ^{In}	λ^{LS}	λ^{In}
R_{\min}^2 or ℓ_{\max}	0.99995	0.06	0.9999	0.06	0.9999	0.06
Number of samples	20	56	140	62	90	64
Min.	-1.666	-1.682	-1.692	-1.677	-1.667	-1.673
1st Qu.	-1.663	-1.677	-1.676	-1.671	-1.663	-1.666
Median	-1.662	-1.672	-1.671	-1.668	-1.658	-1.657
Mean	-1.661	-1.672	-1.671	-1.667	-1.658	-1.658
3rd Qu.	-1.659	-1.666	-1.667	-1.665	-1.653	-1.650
Max	-1.654	-1.662	-1.651	-1.657	-1.646	-1.645

where the density of θ is $\frac{1}{2} \sin(\theta) \mathbf{1}_{[0, \pi]}(\theta)$ and where φ is uniform on $[0, 2\pi)$. The simulation of θ is achieved using the standard method by $2 \arcsin(\sqrt{U})$ where U is uniform on $[0, 1)$. The distribution function of the exit time from the unit ball is obtained via Monte Carlo simulations.

We note that once again we obtain an accuracy of about three digits on the principal eigenvalue using either the WOR or the WOS method. The two statistical methods lead to very similar approximations. If we look at the CPU times, we observe that they increase only linearly with the dimension of the problem.

5.2. A 5D problem

We now take some more examples in dimension 5 to emphasize that our method is not very sensitive to the dimensional effect. Similarly to the 3D case, we consider the Dirichlet problem on the domain

$$D_{5D} = \{(x_1, \dots, x_5) \in \mathbb{R}^5; (x_1, x_2) \in D, (x_3, x_4, x_5) \in [0, 4]^3\}.$$

Using again a separation of variables, the first eigenvalue λ_{5D} is $\lambda_{2D} - 3\pi^2/32 \simeq -1.665$, if λ_{2D} is approximated by -0.740 .

In Table 6, we use confidence intervals and least squares approximations for intervals $[t_1, t_2]$ contained in $[2, 6]$ with $t_2 - t_1 \geq 2$ and a step of 0.1.

As the WOS is not so easy to implement in this case, we now use the WOR and the improved Euler scheme in the simulations. We still obtain in less than 10 minutes of CPU a good accuracy of about 3×10^{-3} on the eigenvalue for a problem which is very difficult to solve by means of deterministic methods. The CPU times of the WOR still increases linearly. It seems sufficient to use a discretization parameter of $\Delta t = 10^{-2}$ for the Euler scheme to reach this accuracy. We can certainly recommend this method for such difficult problems as it is very easy to implement.

6. Conclusion

We have presented and tested a Monte Carlo method to compute the first eigenvalue λ_1 of the Laplace operator with Dirichlet boundary condition. This method is based on the numerical computation of the speed of absorption of the Brownian motion by the boundary of the domain. It requires a good approximation of the law of the exit time of the Brownian motion by means of various schemes and also accurate estimators of λ_1 seen as a parameter of a model.

We have to note that the estimators of λ_1 are very sensitive to the quality of the empirical distribution function of the exit time. In our least squares method, we have however developed a test based on the coefficient R^2 , which appears to be robust in all our experiments. The accuracy is anyhow limited by the speed of convergence given by the Kolmogorov–Smirnov theorem.

The WOR-IS is not efficient in our case, while the naive Euler scheme overestimates λ and is also to be rejected for this purpose. Yet some variants of the Euler scheme provide much more better estimates, while remaining easy to set up. The WOS is the fastest method there, but may be difficult to set up in high dimension. For polytopes, the WOR provides a good approximation of λ_1 but takes longer time in high dimension, and is more difficult to implement. Note that however partial tabulations of some random variables and a better use of the WOR-IS should provide very efficient methods.

On all our test cases, from dimensions 2 to 5, we have obtained in no more than 10 min an accuracy of about three digits on the eigenvalue λ_1 using 1,000,000 particles. Of course, in our test cases, the geometry of the domain was rather simple, but the different methods can be combined to deal with more complex geometries. We can also add that our method is even more valuable in dimension greater than 3, where deterministic methods are limited by the difficulty of generating a mesh and the size of the matrix to invert. In all our methods, the computation time increases linearly with the dimension, and none requires the construction of a mesh. Besides, this method can be extended to deal with a general second-order elliptic operator. The computation of λ_1 could also be a good test to evaluate the efficiency of diffusion approximations.

Acknowledgements

Antoine Lejay is grateful to GdR MOMAS (financed by the institutes ANDRA, BRGM, CEA, CNRS and EDF) for its partial support.

References

- [1] A. Borodin, P. Salminen, *Handbook of Brownian Motion: Facts and Formulae*, Birkhäuser, 1996.
- [2] D. Brockway, P. Soran, P. Whalen, Monte Carlo eigenvalue calculation, in: R. Alcouffe, et al. (Eds.), *Monte-Carlo Methods and Applications in Neutronics, Photonics and Statistical Physics*, Springer-Verlag, 1985, pp. 378–384.
- [3] F.M. Buchmann, Simulations of stopped diffusions, *J. Comput. Phys.* 202 (2) (2005) 446–462.
- [4] F. Campillo, A. Lejay, A Monte Carlo method without grid for a fractured porous domain model, *Monte Carlo Meth. Appl.* 8 (2) (2002) 129–148.
- [5] M. Deaconu, A. Lejay, A random walk on rectangles algorithm, *Methodol. Comput. Appl. Probab.* 8 (1) (2006) 135–151.
- [6] M. Deaconu, A. Lejay, Simulation of a diffusion process using the importance sampling paradigm, in preparation.
- [7] M. Freidlin, *Functional Integration and Partial Differential Equations*, Princeton University Press, 1985.
- [8] A. Friedman, *Stochastic Differential Equations and Applications*, vol. 2, Academic Press, 1976.
- [9] E. Gobet, Weak approximations of killed diffusions using Euler schemes, *Stoch. Process. Appl.* 87 (2) (2000) 167–197.
- [10] E. Gobet, Euler schemes and half-space approximations for the simulation of diffusion in a domain, *ESAIM Probab. Stat.* 5 (2001) 261–297.
- [11] E. Gobet, S. Menozzi, Exact approximation rate of killed hypoelliptic diffusions using the discrete Euler scheme, *Stochastic Process. Appl.* 112 (2) (2004) 201–223.
- [12] N. Golyandina, Convergence rate for spherical processes with shifted centers, *Monte Carlo Meth. Appl.* 10 (3–4) (2004) 287–296.
- [13] Y. Kifer, *Random perturbations of dynamical systems*, *Progress in Probability and Statistics*, vol. 16, Birkhäuser, 1988.
- [14] P. Kloeden, E. Platen, *Numerical solution of stochastic differential equations*, *Applications of Mathematics*, Springer-Verlag, Berlin, 1992.
- [15] B. Lapeyre, E. Pardoux, R. Sentis, *Méthodes de Monte-Carlo pour les equations de transport et de diffusion*, Springer-Verlag, 1998.
- [16] S. Maire, Réduction de variance pour l'intégration numérique et pour le calcul critique en transport neutronique, Ph.D. Thesis, Université de Toulon et du Var, 2001.
- [17] S. Maire, D. Talay, On a Monte Carlo method for neutron transport criticality computations. Preprint no. 2005–02 Université de Toulon et du Var, *IMA J. Numer. Anal.* (2006), in press, doi 10.1093/imanum/drl008.
- [18] G. Milstein, M. Tretyakov, Simulation of a space-time bounded diffusion, *Ann. Appl. Probab.* 9 (3) (1999) 732–779.
- [19] R.G. Pinsky, *Positive harmonic functions and diffusion*, *Cambridge Studies in Advanced Mathematics*, vol. 45, Cambridge University Press, 1995.
- [20] K. Sabelfeld, *Monte Carlo Methods in Boundary Value Problems*, Springer, 1991.
- [21] G. Shorack, J. Wellner, *Empirical processes with applications to statistics*, *Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics*, John Wiley & Sons, 1986.